

Package ‘mnt’

September 25, 2026

Type Package

Title Affine Invariant Tests of Multivariate Normality

Version 1.4

Description Various affine invariant multivariate normality tests are provided. It is designed to accompany the survey article Ebner, B. and Henze, N. (2020) <[doi:10.1007/s11749-020-00740-0](https://doi.org/10.1007/s11749-020-00740-0)> titled “Tests for multivariate normality -- a critical review with emphasis on weighted L^2 -statistics”. We implement new and time honoured L^2 -type tests of multivariate normality, such as the Baringhaus-Henze-Epps-Pulley (BHEP) test, the Henze-Zirkler test, the test of Henze-Jiménez-Gamero, the test of Henze-Jiménez-Gamero-Meintanis, the test of Henze-Visage, the Dörr-Ebner-Henze test based on harmonic oscillator and the Dörr-Ebner-Henze test based on a double estimation in a PDE. Secondly, we include the measures of multivariate skewness and kurtosis by Mardia, Koziol, Malkovich and Afifi and Móri, Rohatgi and Székely, as well as the associated tests. Thirdly, we include the tests of multivariate normality by Cox and Small, the ‘energy’ test of Székely and Rizzo, the tests based on spherical harmonics by Mantzoti and Quiroz and the test of Pudalko. All the functions and tests need the data to be a $n \times d$ matrix where n is the sample size (number of rows) and d is the dimension (number of columns).

License CC BY 4.0

Encoding UTF-8

LazyData true

Author Lucas Butsch [aut],
Bruno Ebner [aut, cre],
Jaco Visagie [ctb],
Johann Siemens [ctb]

Imports PearsonDS, Rmpfr, SuppDists, MASS, pracma, utils

Maintainer Bruno Ebner <bruno.ebner@kit.edu>

Depends R (>= 3.5.0)

Config/roxygen2/version 8.1.0

NeedsCompilation no

Repository CRAN

Date/Publication 2026-09-25 17:00:02 UTC

Contents

mnt-package	3
BHEP	4
CS	5
cv.quan	6
DEHT	7
DEHU	8
EHS	8
HJG	9
HJM	10
HV	11
HZ	12
KKurt	13
MAKurt	14
MASkew	15
MKurt	16
MQ1	17
MQ2	17
MRSSkew	18
MSkew	19
print.mnt	20
PU	20
Quantile09	21
Quantile095	22
Quantile099	22
SR	23
standard	24
test.BHEP	24
test.CS	26
test.DEHT	28
test.DEHU	29
test.EHS	30
test.HJG	31
test.HJM	32
test.HV	33
test.HZ	34
test.KKurt	36
test.MAKurt	37
test.MASkew	38
test.MKurt	40
test.MQ1	41
test.MQ2	42
test.MRSSkew	43
test.MSkew	44
test.PU	45
test.SR	46

Description

Various affine invariant multivariate normality tests are provided. It is designed to accompany the survey article Ebner, B. and Henze, N. (2020) [doi:10.1007/s11749020007400](https://doi.org/10.1007/s11749020007400) titled "Tests for multivariate normality – a critical review with emphasis on weighted L^2 -statistics". We implement new and time honoured L^2 -type tests of multivariate normality, such as the Baringhaus-Henze-Epps-Pulley (BHEP) test, the Henze-Zirkler test, the test of Henze-Jiménez-Gamero, the test of Henze-Jiménez-Gamero-Meintanis, the test of Henze-Visage, the Dörr-Ebner-Henze test based on harmonic oscillator and the Dörr-Ebner-Henze test based on a double estimation in a PDE. Secondly, we include the measures of multivariate skewness and kurtosis by Mardia, Koziol, Malkovich and Afifi and Móri, Rohatgi and Székely, as well as the associated tests. Thirdly, we include the tests of multivariate normality by Cox and Small, the 'energy' test of Székely and Rizzo, the tests based on spherical harmonics by Manzotti and Quiroz and the test of Pudalko. All the functions and tests need the data to be a $n \times d$ matrix where n is the sample size (number of rows) and d is the dimension (number of columns).

Use with model residuals

The tests implemented in `mnt` are calibrated for independent and identically distributed observations. Applying them to residuals from fitted models requires additional justification, because parameter estimation may change the null distribution and invalidate the usual critical values. As discussed by Jimenez-Gamero (2020) and acknowledged by Ebner and Henze (2020), validity depends on both the test and the underlying model.

Author(s)

Maintainer: Bruno Ebner <bruno.ebner@kit.edu>

Authors:

- Bruno Ebner <bruno.ebner@kit.edu>
- Lucas Butsch <lucas.butsch@gmail.com>

Other contributors:

- Jaco Visagie <jaco.visagie@nwu.ac.za> [contributor]
- Johann Siemens <johann-siemens@web.de> [contributor]

References

- Jimenez-Gamero, M. D. (2020). Comments on: Tests for multivariate normality—a critical review with emphasis on weighted L^2 -statistics. *TEST*, 29, 893–897. [doi:10.1007/s1174902000742y](https://doi.org/10.1007/s1174902000742y).
- Ebner, B. and Henze, N. (2020). Rejoinder on: Tests for multivariate normality—a critical review with emphasis on weighted L^2 -statistics. *TEST*, 29, 911–913. [doi:10.1007/s1174902000744w](https://doi.org/10.1007/s1174902000744w).

BHEP

*Statistic of the BHEP-test***Description**

This function returns the value of the statistic of the Baringhaus-Henze-Epps-Pulley (BHEP) test as in Henze and Wagner (1997).

Usage

```
BHEP(data, a = 1)
```

Arguments

`data` a $n \times d$ matrix of d dimensional data vectors.
`a` positive numeric number (tuning parameter).

Details

The test statistic is

$$BHEP_{n,\beta} = \frac{1}{n} \sum_{j,k=1}^n \exp\left(-\frac{\beta^2 \|Y_{n,j} - Y_{n,k}\|^2}{2}\right) - \frac{2}{(1 + \beta^2)^{d/2}} \sum_{j=1}^n \exp\left(-\frac{\beta^2 \|Y_{n,j}\|^2}{2(1 + \beta^2)}\right) + \frac{n}{(1 + 2\beta^2)^{d/2}}.$$

Here, $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $j = 1, \dots, n$, are the scaled residuals, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error.

Value

value of the test statistic.

References

Henze, N., and Wagner, T. (1997), A new approach to the class of BHEP tests for multivariate normality, *J. Multiv. Anal.*, 62:1–23, [doi:10.1006/jmva.1997.1684](https://doi.org/10.1006/jmva.1997.1684)
Epps T.W., Pulley L.B. (1983), A test for normality based on the empirical characteristic function, *Biometrika*, 70:723–726, [doi:10.1093/biomet/70.3.723](https://doi.org/10.1093/biomet/70.3.723)

Examples

```
BHEP(MASS::mvrnorm(50, c(0, 1), diag(1, 2)))
```

CS *Statistic of the test of Cox and Small*

Description

This function returns the (approximated) value of the test statistic of the test of Cox and Small (1978).

Usage

```
CS(data, Points = NULL)
```

Arguments

data a n x d matrix of d dimensional data vectors.

Points points for approximation of the maximum on the sphere. Points=NULL generates 5000 uniformly distributed Points on the d dimensional unit sphere.

Details

The test statistic is $T_{n,CS} = \max_{b \in \{x \in \mathbf{R}^d: \|x\|=1\}} \eta_n^2(b)$, where

$$\eta_n^2(b) = \frac{\left\| n^{-1} \sum_{j=1}^n Y_{n,j} (b^\top Y_{n,j})^2 \right\|^2 - \left(n^{-1} \sum_{j=1}^n (b^\top Y_{n,j})^3 \right)^2}{n^{-1} \sum_{j=1}^n (b^\top Y_{n,j})^4 - 1 - \left(n^{-1} \sum_{j=1}^n (b^\top Y_{n,j})^3 \right)^2}$$

. Here, $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $j = 1, \dots, n$, are the scaled residuals, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error. Note that the maximum functional has to be approximated by a discrete version, for details see Ebner (2012).

Value

approximation of the value of the test statistic of the test of Cox and Small (1978).

References

Cox, D.R. and Small, N.J.H. (1978), Testing multivariate normality, *Biometrika*, 65:263–272.

Ebner, B. (2012), Asymptotic theory for the test for multivariate normality by Cox and Small, *Journal of Multivariate Analysis*, 111:368–379.

Examples

```
CS(MASS::mvrnorm(50, c(0, 1), diag(1, 2)))
```

`cv.quan`*Monte Carlo simulation of quantiles for normality tests*

Description

This function returns the quantiles of a test statistic with optional tuning parameter.

Usage

```
cv.quan(  
  samplesize,  
  dimension,  
  quantile,  
  statistic,  
  tuning = NULL,  
  repetitions = 1e+05  
)
```

Arguments

<code>samplesize</code>	samplesize for which the empirical quantile should be calculated.
<code>dimension</code>	a natural number to specify the dimension of the multivariate normal distribution
<code>quantile</code>	a number between 0 and 1 to specify the quantile of the empirical distribution of the considered test
<code>statistic</code>	a function specifying the test statistic.
<code>tuning</code>	the tuning parameter of the test statistic.
<code>repetitions</code>	number of Monte Carlo runs.

Value

empirical quantile of the test statistic.

Examples

```
cv.quan(samplesize=10, dimension=2, quantile=0.95, statistic=BHEP, tuning=2.5, repetitions=1000)
```

DEHT

*Statistic of the DEH test based on harmonic oscillator***Description**

Computes the test statistic of the DEH test.

Usage

```
DEHT(data, a = 1)
```

Arguments

`data` a $n \times d$ numeric matrix of data values.
`a` positive numeric number (tuning parameter).

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a . Each row of the data Matrix contains one of the n (multivariate) sample with dimension d . To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

The value of the test statistic.

References

Dörr, P., Ebner, B., Henze, N. (2020) "Testing multivariate normality by zeros of the harmonic oscillator in characteristic function spaces", Scandinavian Journal of Statistics, 48(2): 456-501, [doi:10.1111/sjos.12477](https://doi.org/10.1111/sjos.12477).

Examples

```
DEHT(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=1)
```

DEHU

*Statistic of the DEH test based on a double estimation in PDE***Description**

Computes the test statistic of the DEH based on a double estimation in PDE test.

Usage

```
DEHU(data, a)
```

Arguments

data	a (d,n) numeric matrix containing the data.
a	positive numeric number (tuning parameter).

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a. Each row of the data Matrix contains one of the n (multivariate) sample with dimension d. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

The value of the test statistic.

References

Dörr, P., Ebner, B., Henze, N. (2021) "A new test of multivariate normality by a double estimation in a characterizing PDE" *Metrika* 84: 401-427, [doi:10.1007/s0018402000795x](https://doi.org/10.1007/s0018402000795x).

EHS

*Statistic of the EHS test based on a multivariate Stein equation***Description**

Computes the test statistic of the EHS test based on a multivariate Stein equation.

Usage

```
EHS(data, a = 1)
```

Arguments

data	a (d,n) numeric matrix containing the data.
a	positive numeric number (tuning parameter).

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a . Each row of the data Matrix contains one of the n (multivariate) sample with dimension d . To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Note that $a=\text{Inf}$ returns the limiting test statistic with value $2*\text{MSkew} + \text{MRSSkew}$ and $a=0$ returns the value of the limit statistic

$$T_{n,0} = \frac{d}{2} - 2^{\frac{d}{2}+1} \frac{1}{n} \sum_{j=1}^n \|Y_{n,j}\|^2 \exp\left(-\frac{\|Y_{n,j}\|^2}{2}\right).$$

Value

The value of the test statistic.

References

Ebner, B., Henze, N., Strieder, D. (2022) "Testing normality in any dimension by Fourier methods in a multivariate Stein equation", The Canadian Journal of Statistics, 50(3): 992-1033, doi:[10.1002/cjs.11670](https://doi.org/10.1002/cjs.11670).

HJG

Henze-Jiménes-Gamero test statistic

Description

Computes the test statistic of the Henze-Jimenes-Gamero test.

Usage

```
HJG(data, a = 5)
```

Arguments

data	a n x d numeric matrix of data values.
a	positive numeric number (tuning parameter).

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a . Each row of the data Matrix contains one of the n (multivariate) sample with dimension d . To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error.

Value

The value of the test statistic.

References

Henze, N., Jiménez-Gamero, M.D. (2019) "A new class of tests for multinormality with i.i.d. and garch data based on the empirical moment generating function", TEST, 28, 499-521, [doi:10.1007/s117490180589z](https://doi.org/10.1007/s117490180589z)

Examples

```
HJG(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=5)
```

HJM

statistic of the Henze-Jiménez-Gamero-Meintanis test

Description

Computes the test statistic of the Henze-Jiménez-Gamero-Meintanis test.

Usage

```
HJM(data, a)
```

Arguments

data a n x d numeric matrix of data values.
a positive numeric number (tuning parameter).

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a. Each row of the data Matrix contains one of the n (multivariate) sample with dimension d. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error.

Value

The value of the test statistic.

References

Henze, N., Jiménez-Gamero, M.D., Meintanis, S.G. (2019), Characterizations of multinormality and corresponding tests of fit, including for GARCH models, Econometric Th., 35:510–546, [doi:10.1017/S0266466618000154](https://doi.org/10.1017/S0266466618000154).

Examples

```
HJM(MASS::mvrnorm(20,c(0,1),diag(1,2)),a=2.5)
```

HV	<i>statistic of the Henze-Visagie test</i>
----	--

Description

Computes the test statistic of the Henze-Visagie test.

Usage

```
HV(data, a = 5)
```

Arguments

data	a n x d numeric matrix of data values.
a	numeric number greater than 1 (tuning parameter).

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a. Each row of the data Matrix contains one of the n (multivariate) sample with dimension d. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error.

Note that a=Inf returns the limiting test statistic with value $2*MSkew + MRSSkew$.

Value

The value of the test statistic.

References

Henze, N., Visagie, J. (2019) "Testing for normality in any dimension based on a partial differential equation involving the moment generating function", Ann. Inst. Stat. Math., 72:1109–1136, [doi:10.1007/s10463019007208](https://doi.org/10.1007/s10463019007208)

Examples

```
HV(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=5)
HV(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=Inf)
```

 HZ

Statistic of the Henze-Zirkler test

Description

This function returns the value of the statistic of the [BHEP](#) test as in Henze and Zirkler (1990). The difference to the [BHEP](#) test is in the choice of the tuning parameter β .

Usage

```
HZ(data)
```

Arguments

`data` a $n \times d$ matrix of d dimensional data vectors.

Details

A [BHEP](#) test is performed with tuning parameter β chosen in dependence of the sample size n and the dimension d , namely

$$\beta = \frac{((2d + 1)n/4)^{1/(d + 4)}}{\sqrt{2}}.$$

Value

value of the test statistic.

References

Henze, N., and Zirkler, B. (1990), A class of invariant consistent tests for multivariate normality, Commun.-Statist. – Th. Meth., 19:3595–3617, doi:[10.1080/03610929008830400](https://doi.org/10.1080/03610929008830400)

See Also

[BHEP](#)

Examples

```
HZ(MASS::mvrnorm(50,c(0,1),diag(1,2)))
```

KKurt

Koziols measure of multivariate sample kurtosis

Description

This function computes the invariant measure of multivariate sample kurtosis due to Koziol (1989).

Usage

```
KKurt(data)
```

Arguments

data a n x d matrix of d dimensional data vectors.

Details

Multivariate sample kurtosis due to Koziol (1989) is defined by

$$\tilde{b}_{n,d}^{(2)} = \frac{1}{n^2} \sum_{j,k=1}^n (Y_{n,j}^\top Y_{n,k})^4,$$

where $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $j = 1, \dots, n$, are the scaled residuals, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error. Note that for $d = 1$, we have a measure proportional to the squared sample kurtosis.

Value

value of sample kurtosis in the sense of Koziol.

References

Koziol, J.A. (1989), A note on measures of multivariate kurtosis, *Biom. J.*, 31:619–624.

Examples

```
KKurt(MASS::mvrnorm(50,c(0,1),diag(1,2)))
```

MAKurt

*multivariate kurtosis in the sense of Malkovich and Afifi***Description**

This function computes the invariant measure of multivariate sample kurtosis due to Malkovich and Afifi (1973).

Usage

```
MAKurt(data, Points = NULL)
```

Arguments

data a n x d matrix of d dimensional data vectors.

Points points for approximation of the maximum on the sphere. Points=NULL generates 1000 uniformly distributed Points on the d dimensional unit sphere.

Details

Multivariate sample skewness due to Malkovich and Afifi (1973) is defined by

$$b_{n,d,M}^{(1)} = \max_{u \in \{x \in \mathbf{R}^d : \|x\|=1\}} \frac{\left(\frac{1}{n} \sum_{j=1}^n (u^\top X_j - u^\top \bar{X}_n)^3 \right)^2}{(u^\top S_n u)^3},$$

where $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error.

Value

value of sample kurtosis in the sense of Malkovich and Afifi.

References

Malkovich, J.F., and Afifi, A.A. (1973), On tests for multivariate normality, J. Amer. Statist. Ass., 68:176–179.

Henze, N. (2002), Invariant tests for multivariate normality: a critical review, Statistical Papers, 43:467–506.

Examples

```
MAKurt(MASS::mvrnorm(50,c(0,1),diag(1,2)))
```

MASkew

*multivariate skewness in the sense of Malkovich and Afifi***Description**

This function computes the invariant measure of multivariate sample skewness due to Malkovich and Afifi (1973).

Usage

```
MASkew(data, Points = NULL)
```

Arguments

data a n x d matrix of d dimensional data vectors.

Points points for approximation of the maximum on the sphere. Points=NULL generates 1000 uniformly distributed Points on the d dimensional unit sphere.

Details

Multivariate sample skewness due to Malkovich and Afifi (1973) is defined by

$$b_{n,d,M}^{(1)} = \max_{u \in \{x \in \mathbf{R}^d: \|x\|=1\}} \frac{\left(\frac{1}{n} \sum_{j=1}^n (u^\top X_j - u^\top \bar{X}_n)^3 \right)^2}{(u^\top S_n u)^3},$$

where $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error.

Value

value of sample skewness in the sense of Malkovich and Afifi.

References

Malkovich, J.F., and Afifi, A.A. (1973), On tests for multivariate normality, J. Amer. Statist. Ass., 68:176–179.

Henze, N. (2002), Invariant tests for multivariate normality: a critical review, Statistical Papers, 43:467–506.

Examples

```
MASkew(MASS::mvrnorm(50,c(0,1),diag(1,2)))
```

MKurt

Mardias measure of multivariate sample kurtosis

Description

This function computes the classical invariant measure of multivariate sample kurtosis due to Mardia (1970).

Usage

```
MKurt(data)
```

Arguments

data a n x d matrix of d dimensional data vectors.

Details

Multivariate sample kurtosis due to Mardia (1970) is defined by

$$b_{n,d}^{(2)} = \frac{1}{n} \sum_{j=1}^n \|Y_{n,j}\|^4,$$

where $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error.

Value

value of sample kurtosis in the sense of Mardia.

References

- Mardia, K.V. (1970), Measures of multivariate skewness and kurtosis with applications, *Biometrika*, 57:519–530.
- Henze, N. (2002), Invariant tests for multivariate normality: a critical review, *Statistical Papers*, 43:467–506.

Examples

```
MKurt(MASS::mvrnorm(50,c(0,1),diag(1,2)))
```

MQ1	<i>first statistic of Manzotti and Quiroz</i>
-----	---

Description

This function returns the value of the first statistic of Manzotti and Quiroz (2001).

Usage

```
MQ1(data)
```

Arguments

data a n x d matrix of d dimensional data vectors.

Value

Value of the test statistic

References

Manzotti, A., and Quiroz, A.J. (2001), Spherical harmonics in quadratic forms for testing multivariate normality, *Test*, 10:87–104, [doi:10.1007/BF02595825](https://doi.org/10.1007/BF02595825)

Examples

```
MQ1(MASS::mvrnorm(50,c(0,1),diag(1,2)))
```

MQ2	<i>second statistic of Manzotti und Quiroz</i>
-----	--

Description

This function returns the value of the second statistic of Manzotti und Quiroz (2001).

Usage

```
MQ2(data)
```

Arguments

data a n x d matrix of d dimensional data vectors.

Value

Value of the test statistic

References

Manzotti, A., and Quiroz, A.J. (2001), Spherical harmonics in quadratic forms for testing multivariate normality, *Test*, 10:87–104, doi:10.1007/BF02595825

Examples

```
MQ2(MASS::mvrnorm(50,c(0,1),diag(1,2)))
```

MRSSkew

multivariate skewness of Móri, Rohatgi and Székely

Description

This function computes the invariant measure of multivariate sample skewness due to Móri, Rohatgi and Székely (1993).

Usage

```
MRSSkew(data)
```

Arguments

`data` a $n \times d$ matrix of d dimensional data vectors.

Details

Multivariate sample skewness due to Móri, Rohatgi and Székely (1993) is defined by

$$\tilde{b}_{n,d}^{(1)} = \frac{1}{n} \sum_{j=1}^n \|Y_{n,j}\|^2 \|Y_{n,k}\|^2 Y_{n,j}^\top Y_{n,k},$$

where $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error. Note that for $d = 1$, it is equivalent to skewness in the sense of Mardia.

Value

value of sample skewness in the sense of Móri, Rohatgi and Székely.

References

Móri, T. F., Rohatgi, V. K., Székely, G. J. (1993), On multivariate skewness and kurtosis, *Theory of Probability and its Applications*, 38:547–551.

Henze, N. (2002), Invariant tests for multivariate normality: a critical review, *Statistical Papers*, 43:467–506.

MSkew

*Mardias measure of multivariate sample skewness***Description**

This function computes the classical invariant measure of multivariate sample skewness due to Mardia (1970).

Usage

```
MSkew(data)
```

Arguments

data a n x d matrix of d dimensional data vectors.

Details

Multivariate sample skewness due to Mardia (1970) is defined by

$$b_{n,d}^{(1)} = \frac{1}{n^2} \sum_{j,k=1}^n (Y_{n,j}^\top Y_{n,k})^3,$$

where $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the function returns an error. Note that for $d = 1$, we have a measure proportional to the squared sample skewness.

Value

value of sample skewness in the sense of Mardia.

References

Mardia, K.V. (1970), Measures of multivariate skewness and kurtosis with applications, *Biometrika*, 57:519–530.

Henze, N. (2002), Invariant tests for multivariate normality: a critical review, *Statistical Papers*, 43:467–506.

Examples

```
MSkew(MASS::mvrnorm(50,c(0,1),diag(1,2)))
```

print.mnt	<i>Print method for tests of multivariate normality</i>
-----------	---

Description

Printing objects of class "mnt".

Usage

```
## S3 method for class 'mnt'
print(x, ...)
```

Arguments

x	object of class "mnt".
...	further arguments to be passed to or from methods.

Details

A mnt object is a named list of numbers and character string, supplemented with test (the name of the teststatistic). test is displayed as a title. The remaining elements are given in an aligned "name = value" format.

Value

the argument x, invisibly, as for all [print](#) methods.

Examples

```
print(test.DEHU(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=1,MC=500))
```

PU	<i>Statistic of the Pudelko test</i>
----	--------------------------------------

Description

Approximates the test statistic of the Pudelko test.

Usage

```
PU(data, r = 2)
```

Arguments

data	a n x d numeric matrix of data values.
r	a positive number (radius of Ball)

Details

This functions evaluates the test statistic with the given data and the specified parameter r . Since since one has to calculate the supremum of a function inside a d -dimensional Ball of radius r . In this implementation the `optim` function is used.

Value

approximate Value of the test statistic

References

Pudelko, J. (2005), On a new affine invariant and consistent test for multivariate normality, Probab. Math. Statist., 25:43–54.

Examples

```
PU(MASS::mvrnorm(20,c(0,1),diag(1,2)),r=2)
```

Quantile09	<i>Simulated empirical 90% quantiles of the tests contained in package mnt</i>
------------	--

Description

A dataset containing the empirical 0.9 quantiles of the tests for the dimensions $d=2, 3, 5$ and samplesizes $n=20, 50, 100$ based on a Monte Carlo Simulation study with 100000 repetitions. The following parameters were used:

- For `BHEP` the parameter $a=1$,
- for `HV` the parameter $a=5$,
- for `HJG` the parameter $a=1.5$,
- for `HJM` the parameter $a=1.5$,
- for `DEHT` the parameter $a=0.25$,
- for `DEHU` the parameter $a=0.5$,
- for `CS` the parameter `Points=NULL`,
- for `PU` the parameter $r=2$,
- for `MASkew` the parameter `Points=NULL`,
- for `MAKurt` the parameter `Points=NULL`,

Usage

```
Quantile09
```

Format

A data frame with 9 rows and 20 columns.

Quantile095	<i>Simulated empirical 95% quantiles of the tests contained in package mnt</i>
-------------	--

Description

A dataset containing the empirical 0.95 quantiles of the tests for the dimensions $d=2, 3, 5$ and samplesizes $n=20, 50, 100$ based on a Monte Carlo Simulation study with 100000 repetitions. The following parameters were used:

- For [BHEP](#) the parameter $a=1$,
- for [HV](#) the parameter $a=5$,
- for [HJG](#) the parameter $a=1.5$,
- for [HJM](#) the parameter $a=1.5$,
- for [DEHT](#) the parameter $a=0.25$,
- for [DEHU](#) the parameter $a=0.5$,
- for [CS](#) the parameter $\text{Points}=\text{NULL}$,
- for [PU](#) the parameter $r=2$,
- for [MASkew](#) the parameter $\text{Points}=\text{NULL}$,
- for [MAKurt](#) the parameter $\text{Points}=\text{NULL}$,

Usage

Quantile095

Format

A data frame with 9 rows and 20 columns.

Quantile099	<i>Simulated empirical 99% quantiles of the tests contained in package mnt</i>
-------------	--

Description

A dataset containing the empirical 0.99 quantiles of the tests for the dimensions $d=2, 3, 5$ and samplesizes $n=20, 50, 100$ based on a Monte Carlo Simulation study with 100000 repetitions. The following parameters were used:

- For [BHEP](#) the parameter $a=1$,
- for [HV](#) the parameter $a=5$,
- for [HJG](#) the parameter $a=1.5$,

- for [HJM](#) the parameter $a=1.5$,
- for [DEHT](#) the parameter $a=0.25$,
- for [DEHU](#) the parameter $a=0.5$,
- for [CS](#) the parameter `Points=NULL`,
- for [PU](#) the parameter $r=2$,
- for [MASkew](#) the parameter `Points=NULL`,
- for [MAKurt](#) the parameter `Points=NULL`,

Usage

```
Quantile099
```

Format

A data frame with 9 rows and 20 columns.

SR	<i>statistic of the Székely-Rizzo test</i>
----	--

Description

This function returns the value of the statistic of the test of multivariate normality (also called *energy test*) as in Székely and Rizzo (2005). Note that the scaled residuals use another scaling in the estimator of the covariance matrix as the other functions of the package `mnt`! It is equivalent to the function [mvnorm.e](#).

Usage

```
SR(data, abb = 1e-08)
```

Arguments

<code>data</code>	a $n \times d$ matrix of d dimensional data vectors.
<code>abb</code>	Stop criterium.

Value

value of the test statistic.

References

Székely, G., and Rizzo, M. (2005), A new test for multivariate normality, *J. Multiv. Anal.*, 93:58–80, [doi:10.1016/j.jmva.2003.12.002](https://doi.org/10.1016/j.jmva.2003.12.002)

See Also

[mvnorm.e](#)

Examples

```
SR(MASS: :mvrnorm(50,c(0,1),diag(1,2)))
```

standard	<i>Empirical scaled residuals</i>
----------	-----------------------------------

Description

A function that computes the scaled residuals of the data.

Usage

```
standard(data)
```

Arguments

data a n x d matrix of d dimensional data vectors..

Value

A n x d matrix of the scaled residuals.

test.BHEP	<i>Baringhaus-Henze-Epps-Pulley (BHEP) test</i>
-----------	---

Description

Performs the BHEP test of multivariate normality of Henze and Wagner (1997), with tuning parameter α . Critical values and p-values are obtained by Monte Carlo simulation or by fitting a Pearson or Johnson distribution to the first four moments of the limiting null distribution.

Usage

```
test.BHEP(
  data,
  a = 1,
  MC.rep = 10000,
  alpha = 0.05,
  cv.method = c("MC", "Pearson", "Johnson")
)
```

Arguments

data	A numeric $n \times d$ matrix with observations in rows. A numeric vector is treated as a univariate sample. All entries must be finite, $n \geq d + 1$, and the sample covariance must be nonsingular.
a	Positive, finite tuning parameter; corresponds to β in the formulas below.
MC.rep	Positive integer giving the number of Monte Carlo replications. Used only when <code>cv.method = "MC"</code> .
alpha	Significance level, strictly between zero and one.
cv.method	Method used to approximate the critical value: "MC" (default), "Pearson", or "Johnson". Partial matching is supported.

Details

The test statistic is

$$BHEP_{n,\beta} = \frac{1}{n} \sum_{j,k=1}^n \exp\left(-\frac{\beta^2 \|Y_{n,j} - Y_{n,k}\|^2}{2}\right) - \frac{2}{(1 + \beta^2)^{d/2}} \sum_{j=1}^n \exp\left(-\frac{\beta^2 \|Y_{n,j}\|^2}{2(1 + \beta^2)}\right) + \frac{n}{(1 + 2\beta^2)^{d/2}}.$$

Here $\beta = a$, $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, and $S_n = n^{-1} \sum_{j=1}^n (X_j - \bar{X}_n)(X_j - \bar{X}_n)^\top$.

With `cv.method = "MC"`, `cv.quan` simulates the null distribution at the observed sample size n and dimension d , using `MC.rep` replications. The same simulated statistics are used for the critical value and the Monte Carlo p-value

$$p_{MC} = \frac{1 + \sum_{b=1}^B \mathbf{1}\{T_b \geq T_{obs}\}}{B + 1},$$

where B is `MC.rep`. The critical value retains the order-statistic convention of `cv.quan`, and Decision remains `Test.value > cv`. Because the Monte Carlo p-value is discrete and uses the plus-one correction, this decision can differ from `pvalue <= alpha` near the cutoff.

With `cv.method = "Pearson"` or `"Johnson"`, the critical value is obtained from a moment-based approximation to the limiting null distribution $T_\beta(d)$. Let κ_r denote its cumulants. The first three cumulants are given by Henze and Wagner (1997), and the fourth by Theorem 3.2 and Appendix A of Ebner and Henze's announced manuscript. The mean, variance, skewness and ordinary kurtosis are

$$\mu = \kappa_1, \quad \sigma^2 = \kappa_2, \quad \gamma_1 = \kappa_3/\kappa_2^{3/2}, \quad \beta_2 = 3 + \kappa_4/\kappa_2^2.$$

In particular, the fourth central moment is $\mu_4 = \kappa_4 + 3\kappa_2^2$; it is not the fourth cumulant.

Value

An object of class "mnt", with components:

Test Name of the test.

param Value of the tuning parameter.

Test.value Value of the test statistic.

cv Approximated upper $1 - \alpha$ critical value.

Decision TRUE if `Test.value > cv`, so the null hypothesis is rejected at the chosen significance level.

`cv.method` Critical-value method actually used.

`pvalue` Upper-tail p-value computed using the selected method. The Pearson and Johnson p-values are asymptotic approximations.

References

Henze, N. and Wagner, T. (1997). A new approach to the BHEP tests for multivariate normality. *Journal of Multivariate Analysis*, 62, 1–23. doi:[10.1006/jmva.1997.1684](https://doi.org/10.1006/jmva.1997.1684).

Ebner, B. and Henze, N. Four-Moment Approximations and Fast p-Values for BHEP Tests of Multivariate Normality. Supplied manuscript ‘BHEP_fourth_moment.pdf’, Theorem 3.2 and Appendix A.

See Also

[BHEP](#), [cv.quan](#)

Examples

```
set.seed(123)
x <- matrix(rnorm(100), ncol = 2)
test.BHEP(x, MC.rep = 500)
if (requireNamespace("PearsonDS", quietly = TRUE)) {
  test.BHEP(x, cv.method = "Pearson")
}
if (requireNamespace("SuppDists", quietly = TRUE)) {
  test.BHEP(x, cv.method = "Johnson")
}
```

test.CS

multivariate normality test of Cox and Small

Description

Performs the test of multivariate normality of Cox and Small (1978).

Usage

```
test.CS(data, MC.rep = 1000, alpha = 0.05, Points = NULL)
```

Arguments

<code>data</code>	a $n \times d$ matrix of d dimensional data vectors.
<code>MC.rep</code>	number of repetitions for the Monte Carlo simulation of the critical value.
<code>alpha</code>	level of significance of the test.
<code>Points</code>	number of points to approximate the maximum functional on the unit sphere.

Details

The test statistic is $T_{n,CS} = \max_{b \in \{x \in \mathbf{R}^d: \|x\|=1\}} \eta_n^2(b)$, where

$$\eta_n^2(b) = \frac{\left\| n^{-1} \sum_{j=1}^n Y_{n,j} (b^\top Y_{n,j})^2 \right\|^2 - \left(n^{-1} \sum_{j=1}^n (b^\top Y_{n,j})^3 \right)^2}{n^{-1} \sum_{j=1}^n (b^\top Y_{n,j})^4 - 1 - \left(n^{-1} \sum_{j=1}^n (b^\top Y_{n,j})^3 \right)^2}$$

. Here, $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $j = 1, \dots, n$, are the scaled residuals, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error. Note that the maximum functional has to be approximated by a discrete version, for details see Ebner (2012).

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.#'

References

Cox, D.R., Small, N.J.H. (1978), Testing multivariate normality, *Biometrika*, 65:263-272.

Ebner, B. (2012), Asymptotic theory for the test for multivariate normality by Cox and Small, *Journal of Multivariate Analysis*, 111:368-379.

See Also

[CS](#)

Examples

```
test.CS(MASS::mvrnorm(10,c(0,1),diag(1,2)),MC.rep=100)
```

test.DEHT	<i>Doerr-Ebner-Henze test of multivariate normality based on harmonic oscillator</i>
-----------	--

Description

Computes the multivariate normality test of Doerr, Ebner and Henze (2019) based on zeros of the harmonic oscillator.

Usage

```
test.DEHT(data, a = 1, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
a	positive numeric number (tuning parameter).
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value.
alpha	level of significance of the test.

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a. Each row of the data Matrix contains one of the n (multivariate) sample with dimension d. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$param value tuning parameter.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Doerr, P., Ebner, B., Henze, N. (2019) "Testing multivariate normality by zeros of the harmonic oscillator in characteristic function spaces" [arXiv:1909.12624](#)

See Also

[DEHT](#)

Examples

```
test.DEHT(MASS::mvrnorm(20,c(0,1),diag(1,2)),a=1,MC=500)
```

test.DEHU	<i>Doerr-Ebner-Henze test of multivariate normality based on a double estimation in a PDE</i>
-----------	---

Description

Computes the multivariate normality test of Doerr, Ebner and Henze (2019) based on a double estimation in a PDE.

Usage

```
test.DEHU(data, a = 0.5, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
a	positive numeric number (tuning parameter).
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value.
alpha	level of significance of the test.

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a. Each row of the data Matrix contains one of the n (multivariate) sample with dimension d. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$param value tuning parameter.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Doerr, P., Ebner, B., Henze, N. (2019) "Testing multivariate normality by zeros of the harmonic oscillator in characteristic function spaces" [arXiv:1909.12624](#)

See Also[DEHU](#)**Examples**

```
test.DEHU(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=1,MC=500)
```

test.EHS

Ebner-Henze-Strieder test of multivariate normality based on Fourier methods in a multivariate Stein equation

Description

Computes the multivariate normality test of Ebner, Henze and Strieder (2020) based on Fourier methods in a multivariate Stein equation.

Usage

```
test.EHS(data, a = 0.5, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a $n \times d$ matrix of d dimensional data vectors.
a	positive numeric number (tuning parameter).
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value.
alpha	level of significance of the test.

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a . Each row of the data Matrix contains one of the n (multivariate) sample with dimension d . To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level α :

\$Test name of the test.

\$param value tuning parameter.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Ebner, B., Henze, N., Strieder, D. (2020) "Testing normality in any dimension by Fourier methods in a multivariate Stein equation" [arXiv:2007.02596](#)

See Also

[EHS](#)

Examples

```
test.EHS(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=1,MC=500)
```

test.HJG	<i>Henze-Jimenes-Gamero test of multivariate normality</i>
----------	--

Description

Computes the multivariate normality test of Henze and Jimenes-Gamero (2019) in dependence of a tuning parameter a .

Usage

```
test.HJG(data, a = 1, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a $n \times d$ matrix of d dimensional data vectors.
a	positive numeric number (tuning parameter).
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value.
alpha	level of significance of the test.

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a . Each row of the data Matrix contains one of the n (multivariate) sample with dimension d . To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level α :

\$Test name of the test.

\$param value tuning parameter.

```
$Test.value the value of the test statistic.  
$cv the approximated critical value.  
$Decision the comparison of the critical value and the value of the test statistic.#'
```

References

Henze, N., Jimenez-Gamero, M.D. (2019) "A new class of tests for multinormality with i.i.d. and garch data based on the empirical moment generating function", TEST, 28, 499-521, doi:10.1007/s117490180589z

See Also

HJG

Examples

```
test.HJM(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=1.5,MC.rep=500)
```

test.HJM	<i>Henze-Jimenes-Gamero-Meintanis test of multivariate normality</i>
----------	--

Description

Computes the test statistic of the Henze-Jimenes-Gamero-Meintanis test.

Usage

```
test.HJM(data, a = 1.5, MC.rep = 500, alpha = 0.05)
```

Arguments

- data a n x d matrix of d dimensional data vectors.
- a positive numeric number (tuning parameter).
- MC.rep number of repetitions for the Monte Carlo simulation of the critical value.
- alpha level of significance of the test.

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a. Each row of the data Matrix contains one of the n (multivariate) sample with dimension d. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level α :

\$Test name of the test.

\$param value tuning parameter.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Henze, N., Jimenes-Gamero, M.D., Meintanis, S.G. (2019), Characterizations of multinormality and corresponding tests of fit, including for GARCH models, *Econometric Th.*, 35:510-546, doi:[10.1017/S0266466618000154](https://doi.org/10.1017/S0266466618000154).

See Also

[HJM](#)

Examples

```
test.HJM(MASS::mvrnorm(10,c(0,1),diag(1,2)),a=2.5,MC=100)
```

test.HV

The Henze-Visagie test of multivariate normality

Description

Computes the multivariate normality test of Henze and Visagie (2019).

Usage

```
test.HV(data, a = 5, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a $n \times d$ matrix of d dimensional data vectors.
a	positive numeric number (tuning parameter).
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value.
alpha	level of significance of the test.

Details

This functions evaluates the teststatistic with the given data and the specified tuning parameter a . Each row of the data Matrix contains one of the n (multivariate) sample with dimension d . To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Note that $a = \text{Inf}$ returns the limiting test statistic with value $2 * \text{MSkew} + \text{MRSSkew}$.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level α :

`$Test` name of the test.

`$param` value tuning parameter.

`$Test.value` the value of the test statistic.

`$cv` the approximated critical value.

`$Decision` the comparison of the critical value and the value of the test statistic.

References

Henze, N., Visagie, J. (2019) "Testing for normality in any dimension based on a partial differential equation involving the moment generating function", to appear in Ann. Inst. Stat. Math., [doi:10.1007/s10463019007208](https://doi.org/10.1007/s10463019007208)

See Also

[HV](#)

Examples

```
test.HV(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=5,MC.rep=500)
test.HV(MASS::mvrnorm(50,c(0,1),diag(1,2)),a=Inf,MC.rep=500)
```

test.HZ

The Henze-Zirkler test

Description

Performs the test of multivariate normality of Henze and Zirkler (1990).

Usage

```
test.HZ(data, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value
alpha	level of significance of the test

Details

A [BHEP](#) test is performed with tuning parameter β chosen in dependence of the sample size n and the dimension d, namely

$$\beta = \frac{((2d + 1)n/4)^{1/(d + 4)}}{\sqrt{2}}.$$

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$param value tuning parameter.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.#'

References

Henze, N., Zirkler, B. (1990), A class of invariant consistent tests for multivariate normality, Commun.-Statist. - Th. Meth., 19:3595-3617, [doi:10.1080/03610929008830400](#)

See Also

[HZ](#)

Examples

```
test.HZ(MASS::mvrnorm(50,c(0,1),diag(1,2)),MC.rep=500)
```

test.KKurt	<i>Test of normality based on Koziols measure of multivariate sample kurtosis</i>
------------	---

Description

Computes the multivariate normality test based on the invariant measure of multivariate sample kurtosis due to Koziol (1989).

Usage

```
test.KKurt(data, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value
alpha	level of significance of the test

Details

Multivariate sample kurtosis due to Koziol (1989) is defined by

$$\widetilde{b}_{n,d}^{(2)} = \frac{1}{n^2} \sum_{j,k=1}^n (Y_{n,j}^\top Y_{n,k})^4,$$

where $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $j = 1, \dots, n$, are the scaled residuals, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error. Note that for $d = 1$, we have a measure proportional to the squared sample kurtosis.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Koziol, J.A. (1989), A note on measures of multivariate kurtosis, Biom. J., 31:619-624.

See Also[KKurt](#)**Examples**

```
test.KKurt(MASS::mvrnorm(50,c(0,1),diag(1,2)),MC.rep=500)
```

test.MAKurt	<i>Test of normality based on multivariate kurtosis in the sense of Malkovich and Afifi</i>
-------------	---

Description

Computes the multivariate normality test based on the invariant measure of multivariate sample kurtosis due to Malkovich and Afifi (1973).

Usage

```
test.MAKurt(data, MC.rep = 10000, alpha = 0.05, num.points = 1000)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value
alpha	level of significance of the test
num.points	number of points distributed uniformly over the sphere for approximation of the maximum on the sphere.

Details

Multivariate sample skewness due to Malkovich and Afifi (1973) is defined by

$$b_{n,d,M}^{(1)} = \max_{u \in \{x \in \mathbf{R}^d : \|x\|=1\}} \frac{\left(\frac{1}{n} \sum_{j=1}^n (u^\top X_j - u^\top \bar{X}_n)^3 \right)^2}{(u^\top S_n u)^3},$$

where $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level α :

\$Test name of the test.

\$param number of points used in approximation.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Malkovich, J.F., and Afifi, A.A. (1973), On tests for multivariate normality, J. Amer. Statist. Ass., 68:176-179.

Henze, N. (2002), Invariant tests for multivariate normality: a critical review, Statistical Papers, 43:467-506.

See Also

[MAKurt](#)

Examples

```
test.MAKurt(MASS::mvrnorm(10,c(0,1),diag(1,2)),MC.rep=100)
```

test.MASkew	<i>Test of normality based on multivariate skewness in the sense of Malkovich and Afifi</i>
-------------	---

Description

Computes the test of multivariate normality based on skewness in the sense of Malkovich and Afifi (1973).

Usage

```
test.MASkew(data, MC.rep = 10000, alpha = 0.05, num.points = 1000)
```

Arguments

data	a $n \times d$ matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value
alpha	level of significance of the test
num.points	number of points distributed uniformly over the sphere for approximation of the maximum on the sphere.

Details

Multivariate sample skewness due to Malkovich and Afifi (1973) is defined by

$$b_{n,d,M}^{(1)} = \max_{u \in \{x \in \mathbf{R}^d: \|x\|=1\}} \frac{\left(\frac{1}{n} \sum_{j=1}^n (u^\top X_j - u^\top \bar{X}_n)^3 \right)^2}{(u^\top S_n u)^3},$$

where $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$param number of points used in approximation.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Malkovich, J.F., and Afifi, A.A. (1973), On tests for multivariate normality, J. Amer. Statist. Ass., 68:176-179.

Henze, N. (2002), Invariant tests for multivariate normality: a critical review, Statistical Papers, 43:467-506.

See Also

[MASkew](#)

Examples

```
test.MASkew(MASS::mvrnorm(10,c(0,1),diag(1,2)),MC.rep=100)
```

test.MKurt	<i>Test of normality based on Mardias measure of multivariate sample kurtosis</i>
------------	---

Description

Computes the multivariate normality test based on the classical invariant measure of multivariate sample kurtosis due to Mardia (1970).

Usage

```
test.MKurt(data, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value
alpha	level of significance of the test

Details

Multivariate sample kurtosis due to Mardia (1970) is defined by

$$b_{n,d}^{(2)} = \frac{1}{n} \sum_{j=1}^n \|Y_{n,j}\|^4,$$

where $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Mardia, K.V. (1970), Measures of multivariate skewness and kurtosis with applications, *Biometrika*, 57:519-530.

Henze, N. (2002), Invariant tests for multivariate normality: a critical review, *Statistical Papers*, 43:467-506.

See Also[MKurt](#)**Examples**

```
test.MKurt(MASS::mvrnorm(50,c(0,1),diag(1,2)),MC.rep=500)
```

test.MQ1

*Manzotti-Quiroz test 1***Description**

Performs the first test of multivariate normality of Manzotti and Quiroz (2001).

Usage

```
test.MQ1(data, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value
alpha	level of significance of the test

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$param value tuning parameter.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.#'

References

Manzotti, A., Quiroz, A.J. (2001), Spherical harmonics in quadratic forms for testing multivariate normality, *Test*, 10:87-104, doi:[10.1007/BF02595825](https://doi.org/10.1007/BF02595825)

See Also[MQ1](#)

Examples

```
test.MQ1(MASS::mvrnorm(50,c(0,1),diag(1,2)),MC.rep=100)
```

test.MQ2

Manzotti-Quiroz test 2

Description

Performs the second test of multivariate normality of Manzotti and Quiroz (2001).

Usage

```
test.MQ2(data, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value
alpha	level of significance of the test

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.#'

References

Manzotti, A., Quiroz, A.J. (2001), Spherical harmonics in quadratic forms for testing multivariate normality, *Test*, 10:87-104, [doi:10.1007/BF02595825](https://doi.org/10.1007/BF02595825)

See Also

[MQ2](#)

Examples

```
test.MQ2(MASS::mvrnorm(50,c(0,1),diag(1,2)),MC.rep=500)
```

test.MRSSkew	<i>Test of multivariate normality based on the measure of multivariate skewness of Mori, Rohatgi and Szekely</i>
--------------	--

Description

Computes the multivariate normality test based on the invariant measure of multivariate sample skewness due to Mori, Rohatgi and Szekely (1993).

Usage

```
test.MRSSkew(data, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value
alpha	level of significance of the test

Details

Multivariate sample skewness due to Mori, Rohatgi and Szekely (1993) is defined by

$$\tilde{b}_{n,d}^{(1)} = \frac{1}{n} \sum_{j=1}^n \|Y_{n,j}\|^2 \|Y_{n,k}\|^2 Y_{n,j}^\top Y_{n,k},$$

where $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error. Note that for $d = 1$, it is equivalent to skewness in the sense of Mardia.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Mori, T. F., Rohatgi, V. K., Szekely, G. J. (1993), On multivariate skewness and kurtosis, Theory of Probability and its Applications, 38:547-551.

Henze, N. (2002), Invariant tests for multivariate normality: a critical review, Statistical Papers, 43:467-506.

See Also[MRSSkew](#)**Examples**

```
test.MRSSkew(MASS::mvrnorm(50,c(0,1),diag(1,2)),MC.rep=500)
```

test.MSkew	<i>Test of normality based on Mardias measure of multivariate sample skewness</i>
------------	---

Description

Computes the multivariate normality test based on the classical invariant measure of multivariate sample skewness due to Mardia (1970).

Usage

```
test.MSkew(data, MC.rep = 10000, alpha = 0.05)
```

Arguments

data	a n x d matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value
alpha	level of significance of the test

Details

Multivariate sample skewness due to Mardia (1970) is defined by

$$b_{n,d}^{(1)} = \frac{1}{n^2} \sum_{j,k=1}^n (Y_{n,j}^\top Y_{n,k})^3,$$

where $Y_{n,j} = S_n^{-1/2}(X_j - \bar{X}_n)$, $\bar{X}_n$ is the sample mean and S_n is the sample covariance matrix of the random vectors $X_1, \dots, X_n$. To ensure that the computation works properly $n \geq d + 1$ is needed. If that is not the case the test returns an error. Note that for $d = 1$, we have a measure proportional to the squared sample skewness.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level alpha:

\$Test name of the test.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

- Mardia, K.V. (1970), Measures of multivariate skewness and kurtosis with applications, *Biometrika*, 57:519-530.
- Henze, N. (2002), Invariant tests for multivariate normality: a critical review, *Statistical Papers*, 43:467-506.

See Also

[MSkew](#)

Examples

```
test.MSkew(MASS::mvrnorm(50,c(0,1),diag(1,2)),MC.rep=500)
```

test.PU

Pudelko test of multivariate normality

Description

Computes the (approximated) Pudelko test of multivariate normality.

Usage

```
test.PU(data, MC.rep = 10000, alpha = 0.05, r = 2)
```

Arguments

data	a $n \times d$ matrix of d dimensional data vectors.
MC.rep	number of repetitions for the Monte Carlo simulation of the critical value.
alpha	level of significance of the test.
r	a positive number (radius of Ball)

Details

This functions evaluates the test statistic with the given data and the specified parameter r . Since since one has to calculate the supremum of a function inside a d -dimensional Ball of radius r . In this implementation the [optim](#) function is used.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level α :

\$Test name of the test.

\$param value tuning parameter.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.

References

Pudelko, J. (2005), On a new affine invariant and consistent test for multivariate normality, *Probab. Math. Statist.*, 25:43-54.

See Also

[PU](#)

Examples

```
test.PU(MASS::mvrnorm(20,c(0,1),diag(1,2)),r=2,MC=100)
```

test.SR

Szekely-Rizzo (energy) test

Description

Performs the test of multivariate normality of Szekely and Rizzo (2005). Note that the scaled residuals use another scaling in the estimator of the covariance matrix!

Usage

```
test.SR(data, MC.rep = 10000, alpha = 0.05, abb = 1e-08)
```

Arguments

data a $n \times d$ matrix of d dimensional data vectors.

MC.rep number of repetitions for the Monte Carlo simulation of the critical value

alpha level of significance of the test

abb Stop criterium.

Value

a list containing the value of the test statistic, the approximated critical value and a test decision on the significance level α :

\$Test name of the test.

\$Test.value the value of the test statistic.

\$cv the approximated critical value.

\$Decision the comparison of the critical value and the value of the test statistic.#'

References

Szekely, G., Rizzo, M. (2005), A new test for multivariate normality, J. Multiv. Anal., 93:58-80,
[doi:10.1016/j.jmva.2003.12.002](https://doi.org/10.1016/j.jmva.2003.12.002)

See Also

[SR](#)

Examples

```
test.SR(MASS::mvrnorm(50,c(0,1),diag(1,2)),MC.rep=500)
```

Index

* datasets

Quantile09, [21](#)
Quantile095, [22](#)
Quantile099, [22](#)

BHEP, [4](#), [12](#), [21](#), [22](#), [26](#), [35](#)

CS, [5](#), [21–23](#), [27](#)
cv.quan, [6](#), [26](#)

DEHT, [7](#), [21–23](#), [28](#)
DEHU, [8](#), [21–23](#), [30](#)

EHS, [8](#), [31](#)

HJG, [9](#), [21](#), [22](#), [32](#)
HJM, [10](#), [21–23](#), [33](#)
HV, [11](#), [21](#), [22](#), [34](#)
HZ, [12](#), [35](#)

KKurt, [13](#), [37](#)

MAKurt, [14](#), [21–23](#), [38](#)
MASkew, [15](#), [21–23](#), [39](#)
MKurt, [16](#), [41](#)
mnt (mnt-package), [3](#)
mnt-package, [3](#)
MQ1, [17](#), [41](#)
MQ2, [17](#), [42](#)
MRSSkew, [9](#), [11](#), [18](#), [34](#), [44](#)
MSkew, [9](#), [11](#), [19](#), [34](#), [45](#)
mvnorm.e, [23](#)

optim, [21](#), [45](#)

print, [20](#)
print.mnt, [20](#)
PU, [20](#), [21–23](#), [46](#)

Quantile09, [21](#)
Quantile095, [22](#)

Quantile099, [22](#)

SR, [23](#), [47](#)
standard, [24](#)

test.BHEP, [24](#)
test.CS, [26](#)
test.DEHT, [28](#)
test.DEHU, [29](#)
test.EHS, [30](#)
test.HJG, [31](#)
test.HJM, [32](#)
test.HV, [33](#)
test.HZ, [34](#)
test.KKurt, [36](#)
test.MAKurt, [37](#)
test.MASkew, [38](#)
test.MKurt, [40](#)
test.MQ1, [41](#)
test.MQ2, [42](#)
test.MRSSkew, [43](#)
test.MSkew, [44](#)
test.PU, [45](#)
test.SR, [46](#)